

Impact Factor 6.1



# Journal of Cyber Security

ISSN:2096-1146

Scopus

DOI

Google Scholar



More Information

[www.journalcybersecurity.com](http://www.journalcybersecurity.com)



Crossref



Google

Scholar

scopus

# Review Paper on“Fixed Point Theorems in Intuitionistic Fuzzy Metric Space”

1. Arun Kumar Garg 2. Ria Sharma

1. “Professor, Department of Mathematics, UIS, “Chandigarh University, Gharuan, Mohali , Punjab 140413,India”, Email: Arune7246@cumail.in”

2. “Research Scholar,Department of Mathematics, UIS, “Chandigarh University, Gharuan, Mohali, Punjab 140413,India”, Email: riasharma4086@gmail.com”

## **ABSTRACT**

In this paper, as a survey paper, “we evaluate many works related to the fixed point theory for the generalization of fuzzy metric space by using the idea of intuitionistic fuzzy metric space”.

### **1. “INTRODUCTION” :**

Fuzzy set theory was first introduced by Zadeh [1] in 1965 to depict the situation in which data is uncertain. Thereafter the concept of fuzzy set was summed up as intuitionistic fuzzy set by K. Atanassov [2] in 1984. Alaca et al. [3] utilizing the possibility of intuitionistic fuzzy sets, they characterized the notion of intuitionistic fuzzy metric space as Park [4] with the assistance of continuous t-norms and continuous t-conorms. The concept of fuzzy metric space was first introduced by Kramosil and Michalek [5] by utilizing the possibility of Intuitionistic fuzzy set, It has an extensive variety of application in the field of computer programming etc. In 2004, Park [6] characterized the notion of intuitionistic fuzzy metric space with the assistance of continuous t-norm and continuous t-conorm. Al-Thagafi and N. Shahzad [7] demonstrated basic settled point theorems for compatible mappings in complete intuitionistic fuzzy metric spaces. Deng [8], Erceg[9], Kaleva and Seikkala [10] and numerous authors gave the concept of fuzzy metric space in different ways. Grabiec[11] introduced the settled point theory in fuzzy metric space. After that Jungck[8] introduced settled point theorem of intuitionistic fuzzy metric space for commuting mappings. The concept of fuzzy 2-metric space is given by Sushil Sharma[12] in 2002 and he likewise demonstrated basic settled point theorem in fuzzy 2-metric space. They introduced the concept of non Archimedean intuitionistic fuzzy 3 metric spaces by utilizing the concept of Archimedean fuzzy metric space by Dorel Mihet[13], Sushil Sharma[12] and Renu Chugh and Sumitra[14]. Kramosil and Michalek [15] and defined a Hausdorff topology on fuzzy metric space which are often used in current researches. Grabiec [16] extended classical fixed point theorems of Banach and Edelstein to complete and compact fuzzy metric spaces respectively. Park [17] also proved some basic results which include Baire's theorem, separability of the space, second countability of the space and its relation

with 'separability', 'uniform utmost theorem' etc. "Presently, it remains an important problem in fuzzy topology to obtain an appropriate concept of Intuitionistic fuzzy metric spaces". This problem has been investigated by Saadati and Park [18] wherein they defined precompact sets in Intuitionistic fuzzy metric spaces and proved that any subset of an Intuitionistic fuzzy metric space is compact if and just in the event that it is precompact and complete. Also they defined topologically complete Intuitionistic fuzzy metrizable spaces. "The study of common fixed points of mappings satisfying certain contractive conditions has been at the center of strong research activity and, being the area of the fixed point theory, has very important application in applied mathematics and sciences. It was the defining moment in the "fixed point arena" when the idea of commutativity was used by Jungck to obtain a generalization of Banach's fixed point theorem for a pair of mappings". "This result was further generalized and extended in various ways by many authors. Sessa [19] introduced a weaker version of commutativity for a pair of selfmaps, and it is shown that weakly driving pair of maps in metric space is driving, however the converse may not be true". It is well realized that in the setting of metric space, strict contractive condition don't ensure the existence of common fixed point unless the Space is assumed to be compact or the strict conditions are replaced by stronger conditions. "The study of common fixed points of non-compatible mappings is very interesting. Aamri and El Moutawakil [20] generalized the concept of non compatibility in metric spaces by defining the thought property (E.A.) and proved common fixed point theorems under strict contractive conditions. The property (E. An.) allows replacing the completeness requirement of the space with a more natural state of closeness of the range. A major benefit of property (E.A.) is that it ensures. Sharma and Bamoria [21] elucidated a property (SB) in fuzzy metric spaces for self maps".

## 2. "PRELIMINARIES" :

### 2.1 : "SOME BASIC DEFINITIONS USED":

#### Definition1:

"A binary operation,  $\diamond: [0,1] \times [0,1] \rightarrow [0,1]$ , is continuous "t-conorm" if ' $\diamond$ ' is satisfying the following conditions":

- (1) " $\diamond$  is commutative and associative";
- (2) " $\diamond$  is continuous";
- (3) " $a \diamond 0 = a$ ", for all " $a \in [0,1]$ "
- (4) " $a \diamond b \leq c \diamond d$ ", whenever " $a \leq c$ " and " $b \leq d$ " for all " $a, b, c, d \in [0, 1]$ ".

S.NO	AUTHOR	TITLE OF THE PAPER	NAME OF JOURNAL	VOLUME	YEAR	PAGES	MAIN POINTS
1.	“M.Rafi and M.S.M Noorani”	“Fixed point theorem on intuitionistic fuzzy metric space”	“Iranian Journal of Fuzzy Systems”	“3”	“2006”	23-29	“In this paper , authors introduced intuitionistic fuzzy contraction mapping and proved a fixed point theorem in fuzzy metric space.”
2.	“Shams-ur-rahman , M.I. Bhatti, FakharHaider , Muhammad YarBaigand ShabanaAzam”	“Fixed point theorem in intuitionistic fuzzy metric space”	“International Journal of Scientific & Engineering Research”	5	July 2014	1551-1556	“In this paper, authors generalize fuzzy metric space in term of fixed point theorem in Intuitionistic fuzzy metric space.”
3.	“V. Malliga Devi, R. Mohan Raj , and M.Jeyaraman”	“Common Fixed point theorem in intuitionistic fuzzy metric space”	“International Journal of Advanced Engineering Technology”	7	“July-Sept , 2016”	“58-65”	“In this paper, authors proved common fixed point theorems for compatible mappings in complete intuitionistic fuzzy metric”
4.	“NidhiVerma, Dr.RajeshShrivastava”	“A Fixed point theorem in non-archimedean intuitionistic Fuzzy 3-Metric spaces”	“International Journal of Advanced Technology & Engineering Research”	-	2017	112-117	“In this paper,authorsproved a fixed point theorem for a commuting maps and their result generalize and extend some recent results for non Archimedeanintuitionistic fuzzy 3 metric spaces.”

5.	“Akhilesh Jain, Chandel R.S, Hasan Abbas and UdayDolas”	“Common Fixed point theorem in intuitionistic fuzzy metric space under strict contractive conditions”	“International Journal of Recent Scientific Research”	9	July 2018	27716-27721	“In this paper ,authors introduced the notation of Cauchy sequences in an Intuitionistic fuzzy metric space and proved the common fixed point theorem in intuitionistic fuzzy metric space under strict contractive conditions.”
----	---	---	---	---	-----------	-------------	--

**Definition2:**

“Let  $(X, d)$  be a ‘metric space’ and ‘A,B,Sand T’ be four ‘self maps’ on X. ‘The pairs  $(A, S)$  and  $(B, T)$  are said to satisfy common property (E.A.)’, if there exist two sequences  $\{x_n\}$  and  $\{y_n\}$  in X”, such that:

$$\lim_{n \rightarrow \infty} A x_n = \lim_{n \rightarrow \infty} S x_n = \lim_{n \rightarrow \infty} B x_n = \lim_{n \rightarrow \infty} T x_n = t \quad \forall t \in X$$

**Definition3:**

“Let ‘Sand T’ be two ‘self mappings’ of an ‘Intuitionistic fuzzy metric space’  $(X, M, N, *, \diamond)$ , We say that ‘Sand T satisfy the property (SB)’ i.e ( Sharma and Bamboria) if there exists a sequence  $\{x_n\}$  in X” such that:

$$\lim_{n \rightarrow \infty} S x_n = \lim_{n \rightarrow \infty} T x_n = t \quad \forall t \in X$$

**Definition 4:**

“Let A and S be maps from an ‘intuitionistic fuzzy metric space  $(X, M, N, *, \diamond)$ ’ into itself. The maps ‘A and S are said to be weakly commuting’”, if,

$$M(ASz, SAz, t) \geq M(Az, Sz, t) \text{ and } N(Az, Sz, t) \quad \forall z, X \text{ and } t > 0$$

**2.2: ‘LITERATURE SURVEY’ :****2.3: ‘Theorem’ :**

“Let  $(A, B, C, *, \diamond)$  be an ‘intuitionistic fuzzy metric space’ with ‘ $t * t \geq t$ ’ for some ‘ $t \in [0, 1]$ ’ and the condition (FM-6)”. “Let P and Q be ‘weakly compatible mapping’ of A into itself”, such that;

(1) “P and Q satisfy the property (S-B)”,

(2) “there exist a number  $K \in (0,1)$ ” such that ;

$$“B(Qx, Qy, kt) > B(Px, Py, t) * B(Px, Qx, t) * B(Py, Qy, t) * B(Py, Qx, t) * B(Px, Qy, t)”$$

And,

$$“C(Qx, Qy, kt) < C(Px, Py, t) \diamond C(Px, Qx, t) \diamond C(Py, Qy, t) \diamond C(Py, Qx, t) \diamond C(Px, Qy, t)”$$

‘For all  $x \neq y \in A$ ’,

(3) “ $QA \subset PA$  , if  $PA$  or  $QA$  be a close subset of  $A$ , then  $P$  and  $Q$  have a unique common fixed points”.

## 2.4 : ‘Theorem’ :

“Let  $(X, M, N, *, \diamond)$  be an ‘intuitionistic fuzzy metric space’ with ‘ $\iota * \iota \geq \iota$ ’ for some  $\iota \in [0,1]$  and the condition (FM-6)”. “Let  $A, B, S$  and  $T$  be mapping of  $X$  into itself”, such that ;

(1) “ $AX \subset TX$  and  $BX \subset SX$ ”,

(2) “ $(A,S)$  or  $(B,T)$  satisfies the property (S-B)”,

(3) “there exist a number  $k \in (0,1)$ ”, such that;

$$‘M(Ax, By, kt) \geq M(Sx, Ty, t) * M(Sx, By, t) * M(Ty, By, t)’$$

And,

$$‘N(Nx, By, kt) \leq N(Sx, Ty, t) \diamond N(Sx, By, t) \diamond N(Ty, By, t)’ , \text{ for all } x, y \in X,$$

(4) “ $(A,S)$  and  $(B, T)$  are weakly compatible”,

(5) “One of  $AX, BX, SX$ , or  $TX$  is a closed subset of  $X$ ”.

“Then ‘ $A, B, S$  and  $T$ ’ have a unique common fixed point in  $X$ ”.

## 2.5 ‘Theorem’ :

“Let  $(X, M, N, *, \diamond)$  be a ‘complete non Archimedean intuitionistic fuzzy 2 metric space’ and ‘ $q, r: X \rightarrow X$ ’ be a mapping”, satisfying ;

(i) “ $r(X) \subset f(X)$ ”

(ii) “ $r$  is Continuous”

$$(iii) “M(r(x), r(y), a; \alpha t) \geq M(f(x), f(y), a; t) \forall x, y \in X”$$

$X, 0 < \alpha < 1$

And, “ $\lim_{t \rightarrow \infty} M(x, y, a; t) = 1$ ”

‘ $N(r(x), r(y), a; \alpha t) \leq N(f(x), f(y), a; t)$ ’

for all ‘ $x, y \in X$ ’, ‘ $0 < \alpha < 1$ ’

And, ‘ $\lim_{t \rightarrow \infty} N(x, y, a; t) = 0$ ’

‘Then  $q$  and  $r$  have a unique common fixed point’.

## 2.6 ‘Theorem’ :

“Let  $(X, M, N, *, \diamond)$  be a ‘complete non Archimedean intuitionistic fuzzy 3 metric space’ and ‘ $m, n : X \rightarrow X$ ’ be a mapping” satisfying;

(i) “ $n(X) \subset f(X)$ ”

(ii) “ $m$  Continuous”

(iii) “ $M(n(x), n(y), a, b; \alpha t) \geq M(f(x), f(y), a, b; t)$ ”

For all, ‘ $x, y \in X, 0 < \alpha < 1$ ’

And, ‘ $\lim_{t \rightarrow \infty} M(x, y, a, b; t) = 1$ ’

‘ $N(n(x), n(y), a, b; \alpha t) \leq N(m(x), m(y), a, b; t)$ ’

For all  $x, y \in X, 0 < \alpha < 1$

And, ‘ $\lim_{t \rightarrow \infty} N(x, y, a, b; t) = 0$ ’

“Then  $m$  and  $n$  have a unique common point”.

## 3: References:

- 1) C. Alaca, D. Turkoglu, and C. Yildiz, Fixed points in intuitionistic fuzzy metric spaces, Chaos, Solitons and Fractals, 29 (2006), 1073–1078.
- 2) S. Banach, Theorie les operations lineaires, Manograie Matematyczne Warsaw Poland, 1932.
- 3) George and P. Veeramani, on some results in fuzzy metric spaces, Fuzzy Sets and Systems, 64 (1994), no. 3, 395–399.
- 4) J.H. Park, Intuitionistic fuzzy metric spaces, Chaos, Solitons Fractals, 22, No. 5 (2004), 1039-1046, doi:10.1016/j.chaos.2004.02.051
- 5) M. Edelstein, On fixed and periodic points under contractive mappings, Journal of the



- London Mathematical Society, 37 (1962), 74–79
- 6) Park. J.H Intuitionistic fuzzy metric spaces. *Chaos, Solitons& Fractals*, 22:1039:1046, 2004.
  - 7) Al-Thagafi.M.A and N. Shahzad. Generalised I-non expansive self maps and invariants approximations. *Acta Math.Sin.*,24(5):867:876, 2008.
  - 8) Z. K. Deng, Fuzzy pseudo-metric space, *J. Math. Anal.Appl.* 86 (1982), 74-95.
  - 9) M. A. Erceg, Metric space in fuzzy set theory, *J. Math.Anal. Appl.* 69 (1979), 205-230.
  - 10) O. Kaleva and S. Seikkala, On fuzzy metric spaces, *Fuzzy Sets and Systems*, 12 (1984), 215-229.
  - 11) M. Grabiec, Fixed points in fuzzy metric spaces, *Fuzzy Sets and Systems*, (27) (1983), 385-389.
  - 12) Sushil Sharma, On fuzzy metric spaces, *Southeast Asian Bull. Of Math.*, (26) (2002), 133-14
  - 13) Dorel Mihet, Fuzzy  $\psi$  contraction in non-Archimedean fuzzy metric spaces, *Fuzzy Sets and Systems*, (159) (2008),739-744.
  - 14) RenuChugh and Sumitra , Common fixed point theorems in 2 N. A. Menger PM space, *Int. J. Math. Math. Sci.*, ( 26) (8) (2001), 475-483.
  - 15) Kramosil O., Michalek J., Fuzzy metric and statistical metric spaces, *Kybernetika*, 11 (1975), 326-334.
  - 16) Grabiec M., Fixed points in fuzzy metric spaces, *Fuzzy Sets Syst.*27 (1988), 385-389.
  - 17) Park J.S., Kwun Y.C. and Park J.H., A Fixed Point Theorem in the Intuitionistic Fuzzy Metric Spaces, *Far East Journal of Mathematical Sciences*, 16 (2) (2005), 137-149.
  - 18) Saadati R. and Park J. H., On the Intuitionistic Fuzzy Topological Spaces, *Chaos, Solitons and Fractals*, 27(2),2006, 331-344.
  - 19) Sessa S., On weak commutativity condition of mappings in fixed point considerations, *Publ.Inst. Math. Besgrad*,32 (46), (1982) ,149-153.
  - 20) Aamri M., & El Moutawakil, D., Some new common fixed point theorems under strict contractive conditions. *Journal of Mathematical Analysis and Applications*, 270,(2002), 181-188.
  - 21) Sharma Sushil and Bamboria D., Some new common fixed point theorems in fuzzy metric space under strict contractive conditions, *J. Fuzzy Math.*, Vol. 14, No.2(2006), 1-11.