

Impact Factor 6.1



Journal of Cyber Security

ISSN:2096-1146

Scopus

DOI

Google Scholar



More Information

www.journalcybersecurity.com



Crossref



Google

Scholar

scopus

Plant extract: A new class of corrosion inhibitors for low carbon steel in acidic environment

Akhil Saxena

akhil.uis@cumail.in

Chandigarh University Gharuan Mohali

Abstract: By using weight reduction examination, it has been identified that the concentrates of papaya seeds can go about as consumption inhibitor for mellow steel in 15% HCl arrangement. This extract indicates great consumption hindrance proficiency at specific groupings of the corrosive. It is seen that plant removes go about as better inhibitor on expanding their focus.

Keywords: Low carbon steel, inhibitor

Introduction: From most recent couple of decades corrosion inhibitor has been a colorable field of research. Acids are being utilized broadly in the field of gentle steel hardware in enterprises for various applications for e.g. cleaning, descaling, pickling and so on [1]. Various common inhibitors are powerfully used to avoid corrosion and it is the most provident strategy. The hindrance activity of natural inhibitors depends on their adsorption ability on metal by supplanting water atoms [2]. Natural mixes likewise indicate erosion hindrance proficiency and there are a few natural mixes which are informed to forestall consumption [3-7]. In any case, the issue with them is that they are to a great degree noxious to both helpful and additionally environment. In light of the noxious impacts of these natural inhibitors, regular non-harmful inhibitors are the necessity of our condition [1]. Mixes which contain oxygen and nitrogen repress the erosion of gentle steel most strongly [8]. To keep the consumption of mellow steel by utilizing plant extricates is a standout amongst the most venturesome strategies. In green inhibitors auxiliary use O and N are generally present and they are the dynamic places for adsorption [9]. The adsorption of green inhibitors on the steel surface can be either as physisorption or chemisorption or it might likewise be as a consolidated impact of both [10]. Therefore to conquer the poisonous impact of business inhibitors, the advancement of characteristic non-lethal inhibitors to keep the metallic consumption is fundamental and alluring [11]. Here we have discussed the inhibition action of papaya seeds by using weight reduction process.

Experimental:**Weight reduction method:**

The low carbon steel were crushed with fine review emery paper, washed with refined water and dried. At that point the mellow steel strips were weighted precisely before embeddings them into the immersed arrangement of HCl with concentrate. The gentle steel strips were then dunked into HCl without concentrate and soaked with changed convergences of concentrate. At that point we exited the set up for 3 hours and after the slipped by time, the gentle steel strips were rejected, washed down, depleted and gauged. A distinction in weight is noted between the mellow steel strips dunked in HCl arrangement without concentrate and with changed centralizations of concentrate. Here we have utilized 15% HCl in the analysis. Every one of the groupings of inhibitors for weight reduction are taken in mgL⁻¹ by weight. The restraint productivity and surface inclusion (θ) can be dictated by utilizing following condition:

$$\theta = \frac{w_0 - w_i}{w_0} \quad (1)$$

$$\eta(\%) = \frac{w_0 - w_i}{w_0} \times 100 \quad (2)$$

Where, w_i and w_0 are the weight loss values in presence and absence of inhibitor, respectively.

Results and Discussion:

The weight reduction results acquired for mellow steel without and with various convergence of papaya seeds separate in 15% HCl are outlined in Table 1. The table 5 additionally shows that when the centralizations of the extract builds, the hindrance productivity likewise increments. The papaya seeds separate shows most extreme restraint effectiveness of 94% at 300 ppm.

Table 1: Corrosion parameters for mild steel in 15% HCl in absence and presence of different concentrations of *papaya seeds* from weight loss measurements for 3 hours.

Acid solution	Inhibitor concentration(mgL ⁻¹)	Weight loss (mgcm ⁻²)	η (%)	θ
15% HCl	0	36.06	00.00	.0000
	30	07.45	79.34	.7934
	60	06.95	80.73	.8073
	90	05.14	85.74	.8574
	180	04.12	88.57	.8857
	300	02.25	93.76	.9376

Conclusion:

The results indicate that the extract act as an excellent corrosion inhibitor.

References:

1. M. A. Quraishi, A. Singh, V. K. Singh, "Green approach to corrosion inhibition of mild steel in hydrochloric acid and sulphuric acid solutions by the extract of *Murraya koenigii* leaves", *Mater. Chem. Phys.*, vol. 122, page 114-122, 2010.
2. A.S. Fouda, A.S.Ellithy, "Inhibition effect of 4-phenylthiazole derivatives on corrosion of 304L stainless steel in HCl solution", *Corros. Sci.*, vol. 51, page 868, 2009.

3. M. A. Quraishi, R. Sardar, "Dithiazolidines-A new class of heterocyclic inhibitors for prevention of mild steel corrosion in hydrochloric acid solution", *Corros.*, vol. 58, page 103, 2002.
4. H. M. Abd El-Lateef, L. I. Aliyeva, V.M. Abbasov, "Corrosion inhibition of low carbon steel in CO₂-saturated solution using anionic surfactant", *Adv. Appl. Sci.*, vol. 3, page 1185, 2012.
5. M. Lebrini, M. Lagrenee, H. Vezin, L. Gengembre, F. Bentiss, "Electrochemical and quantum chemical studies of new thiadiazole derivatives adsorption on mild steel in normal hydrochloric acid medium", *Corros. Sci.*, vol. 47, page 485, 2005.
6. F. Bentiss, M. Bovanis, B. Mernari, M. Traisnel, M. Legrenee, "Understanding the adsorption of 4H-1, 2, 4-triazole derivatives on mild steel surface in molar hydrochloric acid", *Appl. Surf. Sci.*, vol. 253, page 3696, 2007.
7. G. Husnu, S. H. Ibrahim, "Schinopsis lorentzii as a green corrosion inhibitor for low carbon steel in 1 M HCl", *Ind. Eng. Chem.*, vol. 51, page 785, 2012.
8. D. Yadav, D. Chauhan, I. Hamad, M. A. Quraishi, "Spectroscopic and DFT investigations on the corrosion inhibition behavior of tris (5-methyl-2-thioxo-1, 3, 4-thiadiazole) borate on high carbon steel and aluminium in HCl medium" *RSC Adv.*, vol. 3, page 632, 2013.
9. G. Ji, P. Dwivedi, S. Sundaram, R. Prakash, "Inhibitive effect of chlorophytumborivilianum root extract on mild steel corrosion in HCl and H₂SO₄ solutions", *Ind. Eng. Chem.*, vol. 52, page 10673, 2013.
10. J. Aljourani, K. Raeissi, M.A. Golozar, "Benzimidazole and its derivatives as corrosion inhibitors for mild steel in 1M HCl solution", *Corros. Sci.*, vol. 51, page 1836, 2009.
11. H. Ashassi-Sorkhabi, D. Seifzadeh, M. G. Hosseini, "EN, EIS and polarization studies to evaluate the inhibition effect of 3H-phenothiazin-3-one, 7-dimethylamin on mild steel corrosion in 1M HCl solution", *Corros. Sci.*, vol. 50, page 3363, 2008.